

1 **Prevalence and Determinants of Recent HIV Testing among**
2 **older persons in rural Uganda: A Cross-Sectional Study**

3 Stephen Ojiambo Wandera ^{1, 2*}, Betty Kwagala¹, Fred Maniragaba¹

4 ¹Department of Population Studies, School of Statistics and Planning, College of Business and
5 Management Sciences, Makerere University, Kampala, Uganda

6 ²Demography and Population Studies Programme, Schools of Social Sciences and Public Health,
7 University of the Witwatersrand, South Africa

8 *Corresponding author: swandera@gmail.com

9 Email addresses:

10 BK: kkwagala@gmail.com

11 SOW: swandera@gmail.com

12 FM: fmaniragaba@cartafrica.org

13

14

15

16

17

18

19

20

21

22

23

24

25

26 **Abstract**

27 **Background:** There is limited research on HIV testing among older persons in Uganda. The aim
28 of this study was to investigate the socio-demographic determinants of recent HIV testing among
29 older persons in selected rural districts in Uganda.

30 **Methods:** A cross-sectional survey of 649 older men and women age 50 years and older, from
31 central (Masaka district) and western (Hoima district) Uganda was conducted. Frequency
32 distributions, chi-square tests and multivariable logistic regressions were used to examine the
33 association between recent HIV testing and selected explanatory variables.

34 **Results:** Nearly six in ten (58%) of older persons had primary education. About 60% of the
35 respondents were in union and 13% of them had two or more spouses. Half of the older people
36 (51%) had sex in the last twelve months. A quarter (25%) of older persons gave or received gifts
37 in exchange for sex in their lifetime. Nearly a third (29%) reported sexually transmitted
38 infections in the last 12 months. Prevalence of lifetime HIV testing was 82% and recent (last 12
39 months) HIV testing was 53%. HIV testing in the last 12 months was associated with age
40 (OR=0.50; 95% CI: 0.31-0.79), self-reported sexually transmitted infections (OR=1.59; 95% CI:
41 1.00-2.30), male circumcision (OR=1.71; 95% CI: 1.0-2.93), and sexual activity in the last 12
42 months (OR=2.89; 95% CI: 1.83-4.57).

43 **Conclusion:** Recent HIV testing among older persons was associated with younger age, self-
44 reported STIs, male circumcision, and sexual activity among older persons in rural Uganda. HIV
45 testing interventions need to target older persons who are 70 years and older, who were less
46 likely to test.

47 **Key words:** HIV, AIDS, HIV testing, Aging, STIs, Uganda, Africa

48
49
50
51
52
53
54

55 **Background**

56 The African Union framework on Ageing (AU-Plan) defines older persons as those age 60 years
57 and older [1]. In Uganda, during the drafting of the policy for older persons in 2009, age 60 years
58 and older was used [2, 3]. In this study, we used the World Health Organization (WHO)
59 recommendation of using age 50 and older to define older persons [4]. Subsequently, several
60 studies adopted age 50 and older, as an appropriate definition of old age in sub-Saharan Africa
61 (SSA) countries including Kenya [5, 6]; Uganda [7] and South Africa [8] and those of the WHO
62 Study on global AGEing and adult health (SAGE) and the INDEPTH network [9-14].

63 Globally, the proportion of older persons (age 60 years and older) in 2017 was 13% of the total
64 population [15]. This proportion is projected to increase to 30% by 2050 [16]. Globally, the
65 absolute number of older people increased from 205 million in 1950 to 810 million in 2012 and
66 is anticipated to increase further to 1 billion by 2022 and to 2 billion by 2050 – outnumbering
67 children aged 0-14 years [17-19]. The highest proportion of older persons is projected to live in
68 developing countries by 2050 [20].

69 In Uganda, the proportion of older persons age 50 years and older was estimated at 7.6% in the
70 2014 Uganda Population and Housing Census (UBOS, 2014). There has been some growth in the
71 absolute number of older people in Uganda from 1.1 million in 2002 (4.5% of the population) to
72 1.3 million in 2010 (out of 30 million) and is expected to increase to 5.5 million (constituting
73 5.7% of the population) by 2050 [3, 18].

74 HIV in old age is an emerging public health challenge [21] and considered as a “hidden
75 epidemic” globally [22]. In SSA, only 45% of the general population who are HIV positive know
76 their status [23, 24]. This is far from the UNAIDS ambitious target of ensuring that 90% of those
77 who are HIV positive know their status [25]. With regard to older persons, it is not easy to
78 estimate the prevalence of HIV and the proportion of those who know their HIV status in SSA in
79 general and Uganda in particular [26]. HIV in old age has two major pathways namely; ageing
80 with HIV and infections in old age (Scholten et al., 2011). Ageing with HIV is attributed to
81 adherence to antiretroviral therapy (ART) [27-29].

82 Prevalence of HIV among older persons is estimated at 11% - 13% globally [22, 30]. In sub-
83 Saharan Africa, ten countries (mostly in southern and eastern Africa) account for 80% of all
84 people living with HIV. These include: South Africa (25%), Nigeria (13%), Mozambique (6%),
85 Uganda (6%), Tanzania (6%), Zambia (4%), Zimbabwe (6%), Kenya (6%), Malawi (4%) and
86 Ethiopia (3%) [31-34]

87 HIV testing varies across countries. Among older persons, HIV testing was nearly half (48%) in
88 Uganda [26], 54% in South Africa [35], and 23% in Zimbabwe [36]. However, HIV testing
89 programmes do not prioritize older persons. yet many lack information on HIV prevention and
90 rarely test for HIV/AIDS [37-39]. In addition, older persons experience stigma while accessing
91 HIV testing services [37].

92 The determinants of HIV testing among older persons have been summarized in the conceptual
93 model adapted from the healthcare utilization model. These include predisposing factors (age,
94 gender, race / ethnicity, education, household income, employment status), enabling factors
95 (health insurance, access to care, previous testing, seeing a doctor) and need factors especially
96 HIV risk behaviours in the past 12 months [40-43].

97 Gender is a significant determinant of HIV testing among older persons, where older women are
98 more likely to test for HIV compared to older men [44]. Among older persons, advanced age has
99 been associated with reduced odds of HIV testing [45]. The perceived risk of contracting HIV is
100 an important predictor of HIV testing, where a low perceived risk is associated with reduced
101 odds of HIV testing [46]. Among older persons, prior history of testing reduces the odds of
102 subsequent HIV testing [45].

103 A high level of education increases the odds of HIV testing among older people. With regard to
104 marital status, HIV testing is higher among ever married persons outside union i.e. divorced or
105 separated or widowed [26]. In addition, fear, emotional stress of positive HIV sero status results,
106 HIV stigma [47] and knowledge of HIV transmission [37, 48] have been associated with HIV
107 testing [37].

108 Despite the available evidence on HIV testing, there is dearth of information on HIV testing
109 among older persons in developing countries in general and Uganda [49] in particular. In SSA,

110 sources of data on HIV/AIDS such as Demographic and Health Surveys (DHS) and AIDS
111 Indicator Surveys focus on age 15-54 years. The recent Uganda DHS, 2016 and Population and
112 HIV Impact Surveys (PHIA) in 12 African countries also omit older persons [33, 50] [29]. In
113 addition, several studies on the health of older people in Uganda have focused on later life
114 problems associated with HIV/AIDS (Scholten et al., 2011; Seeley, Wolff, Kabunga,
115 Tumwekwase, & Grosskurth, 2009) but not their uptake of HIV testing. There is need for
116 language editing.

117 Therefore, the aim of this study was to investigate the determinants of access to HIV testing
118 services among older persons in selected rural districts in Uganda. Findings are expected to
119 contribute to understanding of factors associated with HIV testing among older persons [37, 38,
120 45, 51].

121 **Methods**

122 **Study design and setting**

123 The study used a cross-sectional and mixed methods study design. Both survey data and
124 qualitative data were collected in February 2018. Qualitative data included focus group
125 discussions and in-depth interviews. However, this paper is based on survey results. The study
126 setting was in rural districts of Uganda: Masaka (central Uganda) and Hoima (western Uganda).

127 **Study participants**

128 Older persons were defined as those age 50 years and older as recommended by the World
129 Health Organization [4]. We thus considered older persons 50 years and above who had the
130 capacity to provide informed consent.

131 **Sampling procedures**

132 We used a multi-stage stratified cluster sampling design. We randomly selected two regions in
133 Uganda namely: central and western regions out of the four administrative ones. Simple random
134 sampling was used to select one district from each region: Masaka (central) and Hoima
135 (western) from each region. We used the sampling frame of the 2014 Uganda Population and
136 Housing Census [52] to select two sub-counties from Masaka [53] and three sub-counties from

137 Hoima [54] using simple random sampling. From each sub-county, four enumeration areas or
138 villages were selected using simple random sampling. The census sampling frame listed all
139 enumeration areas in each sub-county. Using Microsoft office excel package, computer
140 generated random numbers were assigned to all enumeration areas. Random numbers were used
141 to randomly select enumeration areas. From each village, a sampling frame of older persons' and
142 their households was constructed in consultation with local leaders and systematic sampling was
143 used to select participants for the survey. In households where older men and women live as
144 couples, both were interviewed separately.

145 Kish's formula [55] was applied to generate a sample size of 649 older persons for the survey.
146 The prevalence of HIV testing for those age 50-59 years was 45% among men and 49% among
147 women [56]. We used the lower bound of HIV testing (45%), the $p=0.45$ and the $q=0.55$. The
148 level of confidence was set at 95% ($z=1.96$) and the error at 8% ($e=0.008$). The expected sample
149 size was 148.5. The sample size was multiplied by the design effect of two ($D=2$). Therefore, the
150 expected sample size was 297. The final sample size after adjusting for a response rate of 90%
151 became 330. To allow for small area (district) estimations, the sample size was multiplied by two
152 since the study covered two districts. The overall sample size was 660 older persons. Due to non-
153 response, the final sample size was 649 older persons. The number of older persons selected
154 from each enumeration area was determined by probability proportionate sampling (PPS) from
155 the 2014 Uganda census sampling frame [52].

156 **Data collection procedures**

157 Research assistants were trained for three days by the principal investigator. A pilot study / pre-
158 testing of the tools was conducted in Wakiso town council. Feedback from pilot was integrated
159 in revision of the tools – survey questionnaire and guides for qualitative data. The survey
160 questionnaire was programmed in Android enabled tablets and the SurveyCTO online platform.
161 The survey was conducted in Hoima and Masaka in February 2018. Two research teams
162 (composed of five research assistants) were sent to each district at the same time to collect
163 survey and qualitative data.

164 **Data management using SurveyCTO**

165 Survey data were collected using SurveyCTO [57] application installed on android enabled
166 Tablets. Data were downloaded from the SurveyCTO Server as STATA files on daily basis.

167 **Outcome Variable**

168 The outcome variable was recent HIV testing (last 12 months). Participants were asked if they
169 had ever an HIV test in their life time (yes = 1, no =0) and if they tested for HIV in the last 12
170 months (yes =1 or no = 0). A follow up question was about reception of HIV results during the
171 recent HIV test (yes or no responses). A binary variable called recent HIV testing was recoded to
172 a binary variable (yes and no). Recent HIV testing means HIV testing and reception of results in
173 the last 12 months.

174 **Explanatory variables**

175 Demographic variables included age, and sex. Age was recoded into three categories: 50-59, 60-
176 69 and 70 and older. Sex was recoded into male and female.

177 Socio-economic variables included: education level (none, primary and secondary or higher),
178 working in the last 12 months (yes and no), religion, marital status or currently in union, number
179 of other wives, and children ever born and those currently alive. Religion was recoded as
180 Catholics, Anglicans, Muslims and Others.

181 HIV related variables included knowledge about HIV transmission, HIV stigma, and need to test
182 for HIV (yes or no). These questions were adopted from the Population HIV Impact Assessment
183 (PHIA) survey [33] and the Demographic and Health survey [56]. To measure correct knowledge
184 about HIV transmission, five questions were asked:

- 185 1. Can the risk of HIV transmission be reduced by having sex with only one uninfected
186 partner who has no other partners? (coded as yes =1, no = 0).
- 187 2. A person can get HIV from mosquito bites? (coded as yes =0, no = 1).
- 188 3. Can a person reduce their risk of getting HIV by using a condom every time they have
189 sex? (coded as yes =1, no = 0).
- 190 4. A person cannot get HIV by sharing food with someone who has HIV? (coded as yes =1,
191 no = 0).
- 192 5. Can a healthy-looking person have HIV? (coded as yes =1, no = 0).

193 These questions were recoded as binary variables. Then they were added together to generate an
194 aggregate variable for correct HIV knowledge. The five statements of HIV knowledge had a low
195 reliability test (Cronbach's alpha of 0.43). Correct knowledge about HIV transmission was
196 categorized as agreement to at least four to five statements. Those who had agreement to none to
197 three statements were recoded as not having correct knowledge on HIV transmission.

198 HIV stigma was measured by eight binary (coded as yes = 1 and no = 0) statements (Cronbach's
199 alpha was 0.60):

- 200 1. Would not buy fresh vegetables from an HIV positive vendor
- 201 2. Children living with HIV should not be allowed to attend school with children who do
202 not have HIV
- 203 3. People hesitate to take an HIV test because they are afraid of how other people will react
204 if the test result is positive
- 205 4. People talk badly about people living with HIV, or who are thought to be living with HIV
- 206 5. People living with HIV, or thought to be living with HIV, lose the respect of other people
- 207 6. Fear that one could get HIV if in contact with the saliva of a person living with HIV
- 208 7. Would be ashamed if someone in family had HIV
- 209 8. Not willing to care for someone living with HIV

210 These HIV stigma statements were recoded into binary form and added together to form a score
211 (range from 0 – 8). A binary variable called stigma on at least four statements was created (0 =
212 agreement on 0-3 statements; 1= agreement to 4-8 statements).

213 HIV related behaviour included sexual activity in the last 12 months (yes or no), number of life
214 time sexual partners, transactional sex (life time and recent), alcohol consumption, male
215 circumcision and self-reported STIs. Transactional sex involved giving and receiving of gifts for
216 sex in the last 12 months. Substance use variables included alcohol consumption, smoking and
217 use of tobacco. Males were asked to report about their circumcision status (yes, no and not
218 applicable for females).

219 Self-reported STIs were measured by asking four questions:

- 220 1. During the last 12 months, have you had an abnormal discharge from your vagina or
221 experienced pelvic pain (if woman) or penis (if man)?

- 222 2. During the last 12 months, have you had an ulcer or sore on or near your vagina (woman)
223 or penis (man)?
- 224 3. During the last 12 months, have you had pain on urination?
- 225 4. In the last 12 months, did a doctor, clinical officer or nurse tell you that you had a
226 sexually transmitted disease other than HIV?

227 These questions had three categories (yes, no and don't know). The Cronbach's alpha for the
228 four statements was 0.71. The "don't know" category was merged with the "No" category in
229 order to create binary variables to allow creation of an aggregate variable. In addition, the
230 responses to the former were few. After, an aggregate variable – self-reported STIs was created
231 for those who reported an abnormal discharge, ulcer or sore in the genital area, pain during
232 urination and were told to have an STI by a health provider.

233 **Statistical Analysis**

234 Frequency distributions were used to describe the background characteristics of the older
235 persons. Cross-tabulations were used to investigate associations between recent HIV testing
236 (outcome variable) and selected explanatory variables. Pearson's chi-squared (χ^2) tests were used
237 to examine the significant differences between recent HIV testing and the explanatory variables.
238 The level of statistical significance using p-values was set at $p < 0.05$.

239 Multivariable logistic regression analyses were used to examine the association between recent
240 HIV testing and explanatory variables whose p-values were less than 0.05 during the chi-square
241 tests. We used a step-wise regression for multivariable analysis. The first model includes
242 respondents' background characteristics. In the second model, we added HIV knowledge and
243 attitude factors, and in the third and final model we added behavior factors. Results are presented
244 in the form of Odds Ratios (OR) reporting 95% confidence intervals. The level of statistical
245 significance using p-values was set at $p < 0.05$. All analyses were performed in STATA version
246 15.

247 **Results**

248 **Descriptive characteristics**

249 Table 1 shows the descriptive characteristics of older persons in rural Uganda. About 52% of the
250 respondents were female and 52% were 60 years and older. The majority (75%) had primary or
251 no formal education, were working (53%), either Catholic or Anglican (75%), and were either
252 married or cohabiting (60%).

253 With respect to HIV knowledge and attitude factors, the majority (65%) had correct knowledge
254 on 4-5 HIV transmission statements, had less or no HIV associated stigma (67%), agreed to the
255 need to test for HIV once a year (95%), and the need for male circumcision to prevent HIV
256 (69%). About half (51%) had sex in the past 12 months. Majority (72%) had had one lifetime
257 sexual partner. On the other hand, a small proportion of older people had received or given
258 money or gifts for sex (24%) in their lifetime and in the last 12 months preceding the study
259 (13%), drunk alcohol (34%) and used tobacco or drugs (13%). Some of the respondents reported
260 suffering from STIs in the past 12 months (29%) and were circumcised (17%). Slightly over half
261 (53%) and over three quarters (82%) tested for HIV in the last 12 months and in their lifetime,
262 respectively.

263 **Table 1 about here**

264 **Association between HIV testing in the past 12 months and independent factors**

265 Table 2 shows the association between recent HIV testing and reception of results and
266 background factors among older persons in Uganda.

267 HIV testing prevalence declined with increase in age (64% for 50-59 compared to 19% of 80+
268 year olds; $p<0.001$). Higher proportion of HIV testing were observed among respondents that
269 worked in the past year (59%; $p<0.001$), currently in union (57%; $p<0.01$), polygamous unions
270 (66%; $p<0.01$), had correct knowledge of HIV transmission on 4-5 aspects (56%; $p=0.01$), felt
271 the need to test for HIV annually (54%; $p=0.01$), felt need for male circumcision (56%; $p=0.01$),
272 were actually circumcised (67%; $p<0.01$), had sex in the last 12 months (67%; $p<0.001$), had
273 transactional sex (67%; $p=0.01$) in the last 12 months, and self-reported STIs in the past year

274 (62%; $p < 0.01$). Education, religion, stigma, number of lifetime sexual partners, alcohol and drug
275 use were not significantly associated with HIV testing 12 months preceding the study.

276 **Table 2 about here**

277 **Multivariable results**

278 Table 3 shows the association between recent HIV testing and background factors among older
279 persons in rural Uganda.

280 Age was consistently associated with HIV testing in the past 12 months. Older persons age 70
281 and older consistently had reduced odds of testing for HIV compared to 50-59 year olds after
282 adjusting for background factors (aOR=0.33; 95% CI: 0.22-0.50), adding knowledge factors and
283 attitudes (aOR=0.37; 95% CI: 0.24-0.56) and finally adding behavioural factors (aOR=0.49; 95%
284 CI: 0.31-0.79).

285 In addition, being circumcised, sexual activity, transactional sex, and self-reported STIs in the
286 past year, being circumcised were significantly associated with recent HIV testing. The odds of
287 HIV testing increased (aOR=1.59; 95% CI: 1.0-2.3) among respondents who had a self-reported
288 STI compared to those that did not. Older men who were circumcised compared to those (older
289 men) who were not were more likely (aOR=1.71; 95% CI: 1.0-2.9) to test for HIV in the last 12
290 months. Also, those who had sex in past year compared to those that did not, had increased odds
291 (aOR=2.89; 95% CI: 1.8-4.6) of HIV testing.

292 Working in the past year was significantly associated (aOR=1.39; 95% CI: 1.0-1.9) with HIV
293 testing after adjusting for background characteristics but subsequently lost its influence after
294 adjusting for the rest of the explanatory factors (Models 2 and 3). Likewise, the odds of HIV
295 testing among 60-69 year olds reduced in the first and second models compared to 50-59 year
296 olds but were not significant in the third model. The association between sex, number of spouses,
297 knowledge of HIV transmission modes, stigma, acknowledging the need for circumcision among
298 older person and engaging in transactional sex 12 months prior to the study were not
299 significantly associated with HIV testing 12 months prior to the study.

300 **Table 3 about here**

301 **Discussion**

302 We set out to establish the prevalence and examine the determinants of HIV testing in the last 12
303 months among older persons in rural Uganda. Over half of (53%) older persons had tested HIV
304 during the 12 months preceding the study. This finding is in consistent with the proportion
305 (48%) of older persons (age 45-59 years) that had tested for HIV [37] in the 2011 Uganda AIDS
306 indicator survey [26].

307 The determinants of HIV testing among old persons in the year preceding the study in order of
308 strength of influence were: sexual activity in the past year, male circumcision, having a self-
309 reported STIs and advanced age. Sexual activity in the last 12 months increased the odds of
310 recent HIV testing. Among older persons, recent sexual activity increases perceived risk of HIV
311 infection which motivates older persons to have an HIV test [38, 46]. This is because sexual
312 activity is one of the main avenues of HIV transmission [58]. Older men tend to remain sexually
313 active and engage in extra-marital affairs more than older women [35].

314 Our study found that self-reported STIs were positively associated with HIV testing. Self-
315 reported STIS are indicative of engagement in risky sexual behaviours that increase the odds of
316 HIV infection. HIV testing in such cases is through risk perception or referral by health
317 providers [46].

318 The study found that circumcised older persons had increased odds of HIV testing [59]. The
319 possible explanation is that in case of medical circumcision, it is possible that older persons who
320 interacted with the health sector also benefited from HIV relevant health education [60]. In
321 addition, they might choose not to use condoms and instead prefer to test for HIV.

322 This study found that advanced age reduced odds of testing for HIV. This finding is in
323 agreement with studies elsewhere [45]. This could be associated with low perceived HIV risk,
324 and lack of associated information [46, 61].

325 This study merits the following strengths: first, it highlights important findings about HIV testing
326 among older persons in Uganda using quantitative survey data with a good sample size. The
327 findings provide a benchmark for conducting further studies in Uganda.

328 None the less, there are some limitations of the data. First, it is cross-sectional data and we
329 cannot easily ascertain the direction of causality of associations between HIV testing and self-
330 reported STIs and sexual activity in the last 12 months. Finally, HIV knowledge reliability score
331 of 0.43 showed that the statements are not very reliable.

332 **Conclusion**

333 Recent HIV testing among older persons is associated with younger age (50-59 years), self-
334 reported STIs, male circumcision, and sexual activity among older persons in rural Uganda. HIV
335 testing interventions need to target older persons age 70 years and older, who were less likely to
336 test. These interventions include behavioral risk assessment and routine screening for HIV
337 infection.

338 **Declarations**

339 **Ethics approval and consent to participate**

340 The study protocol was reviewed and approved by The AIDS Support Organization (TASO
341 REC), a local Research and Ethics Committee (REC) on July 4th 2017. The approval reference
342 number is TASOREC/30/17-UG-REC-009. Finally, the protocol was registered with the Uganda
343 National Council of Science and Technology (UNCST), with a registration number, SS 4424.
344 Written voluntary informed consent was obtained from all study participants. Participants were
345 assured of confidentiality. In order to ensure anonymity, participants' names were not be
346 recorded alongside their responses.

347 **Consent for publication**

348 Not applicable

349 **Availability of data and materials**

350 The datasets generated and analysed during the current study are not publicly available due for
351 confidentiality reasons but are available from the corresponding author on reasonable request.

352 **Competing interests**

353 The author(s) declare that they have no competing interests. Stephen Ojiambo Wandera is part of
354 the Editorial board of BMC Public Health, but otherwise, had no role in the editorial process of
355 this article.

356 **Funding**

357 This research was funded by the Consortium for Advanced Research Training in Africa
358 (CARTA) as a Re-Entry grant (RCS/CARTAREENTRY/2016/002). CARTA is jointly led by
359 the African Population and Health Research Center and the University of the Witwatersrand and
360 funded by the Carnegie Corporation of New York (Grant No: B 8606.R02), Sida (Grant No:
361 54100029), the DELTAS Africa Initiative (Grant No: 107768/Z/15/Z). The DELTAS Africa
362 Initiative is an Independent funding scheme of the African Academy of Sciences (AAS)'s
363 Alliance for Accelerating Excellence in Science in Africa (AESA) and supported by the New
364 Partnership for Africa's Development Planning and Coordinating Agency (NEPAD Agency)
365 with funding from the Wellcome Trust (UK) (Grant No: 107768/Z/15/Z) and the UK
366 government.

367 Some seed fund was granted by the Makerere University, College of Business and Management
368 Sciences (CoBAMS) to support manuscript writing process. The funding body (CARTA or
369 Makerere University) had no role in the design of the study; collection, analysis, and
370 interpretation of data; and in the writing the manuscript.

371 **Authors' contributions**

372 SOW conceptualized and developed the study. BK, and FM reviewed the study protocol. BK &
373 FM wrote the background section. BK, PN and SOW reviewed the literature. SOW analysed the
374 data. BK, SOW and SOW interpreted the results and participated in the drafting of the
375 manuscript. All read and reviewed the manuscript. All authors read and approved the manuscript.

376 **Acknowledgements**

377 We acknowledge the statistical support from Ronald Naitala (RN) of Baylor Uganda: Baylor
378 College of Medicine Children's Foundation.

379 **Authors' information**

380 SOW is a Lecturer at the Department of Population Studies (DPS), School of Statistics and
381 Planning (SSP), Makerere University. He has a PhD in Population Studies (specialized on
382 Population Ageing). His PhD thesis was titled: “*Disparities in Health & of Access to Healthcare*
383 *among Older persons in Uganda*”. In addition, he has researched on Sexual and Gender-Based
384 Violence. He holds a Master of Science in Population and Reproductive Health and a Bachelor
385 of Science in Population Studies.

386
387 BK is a Senior Lecturer at the Department of Population Studies, School of Statistics and
388 Planning, College of Business and Management (CoBAMS), Makerere University. BK holds a
389 PhD in Sociology (University of Vienna), Masters in Development Studies (Women and
390 Development), from the Institute of Social Studies (ISS) at The Hague. Her research interests
391 focus on gender and reproductive health.

392 FM is an Assistant Lecturer at the DPS, Makerere University. He holds a Master of Science in
393 Population Studies, Makerere University. He is a PhD student at the Makerere University. He
394 has submitted his thesis for examination. His thesis focused on the “*Determinants of Quality of*
395 *life of older persons in rural Uganda*”.

396 **References**

- 397 1. UNECA: **The state of Older Persons in Africa - 2007**. In: *Regional review and*
398 *appraisal of the Madrid International Plan of Action on Ageing*. United Nations
399 Economic Commission for Africa; 2007.
- 400 2. MoGLSD: **National Policy for Older Persons: Ageing with Security and Dignity** In.
401 Kampala, Uganda: Ministry of Gender, Labour and Social Development; 2009.
- 402 3. UBOS: **Uganda National Household Survey 2009-2010. Socio-economic module.**
403 **Abridged report**. In. Kampala, Uganda: Uganda Bureau of Statistics; 2010.
- 404 4. **Definition of an older or elderly person: Proposed Working Definition of an Older**
405 **Person in Africa for the MDS Project**
406 [<http://www.who.int/healthinfo/survey/ageingdefnolder/en/>]
- 407 5. Kyobutungi C, Ezech A, Zulu E, Falkingham J: **HIV/AIDS and the health of older**
408 **people in the slums of Nairobi, Kenya: results from a cross sectional survey**. *BMC*
409 *Public Health* 2009, **9**(1):153.
- 410 6. Ezech A, Chepngeno-Langat G, Abdhalah Z, Woubalem Z: **The situation of older people**
411 **in poor urban settings: the case of Nairobi, Kenya** In: *Aging in Sub-Saharan Africa:*
412 *Recommendations for Furthering Research* edn. Edited by Cohen B, Menken J.
413 Washington, US: The National Academies Press; 2006: 7-45.

- 414 7. MRC, UVRI: **Direct and indirect effects of HIV/AIDS and anti-retroviral treatment**
415 **on the health and wellbeing of older people.** In: *WHO's Study on global AGEing and*
416 *adult health (SAGE)*. Kampala: Uganda Virus Research Institute & Medical Research
417 Council; 2011.
- 418 8. Gómez-Olivé FX, Thorogood M, Clark B, Kahn K, Tollman S: **Self-reported health and**
419 **health care use in an ageing population in the Agincourt sub-district of rural South**
420 **Africa.** *Global Health Action* 2013, **6**:19305.
- 421 9. HAI: **Guidance on including older people in emergency shelter programmes: a**
422 **summary.** In.: HelpAge International, London, UK 2010.
- 423 10. Kyobutungi C, Egondi T, Ezech A: **The health and well-being of older people in**
424 **Nairobi's slums.** *Global Health Action* 2010, **3**:45-53.
- 425 11. Kowal P, Kahn K, Ng N, Naidoo N, Adbullah S, Bawah A, Binka F, Chuc NTK,
426 Debpuur C, Ezech A *et al*: **Ageing and adult health status in eight lower-income**
427 **countries: the INDEPTH WHO-SAGE collaboration.** *Global Health Action* 2010,
428 **Supplement 2.**
- 429 12. Hirve S, Juvekar S, Lele P, Agarwal D: **Social gradients in self-reported health and**
430 **well-being among adults aged 50 and over in Pune District, India.** *Global Health*
431 *Action* 2010, **2**:88-95.
- 432 13. Gómez-Olivé FX, Thorogood M, Clark B, Kahn K, Tollman S: **Assessing health and**
433 **well-being among older people in rural South Africa.** *Global Health Action* 2010,
434 **2**:23-35.
- 435 14. Debpuur CY, Welaga P, Wak G, Hodgson A: **Self-reported health and functional**
436 **limitations among older people in the Kassena-Nankana District, Ghana; 2010.**
- 437 15. UN: **World Population Prospects: The 2017 Revision, Key Findings and Advance**
438 **Tables** In.: United Nations Department of Economic and Social Affairs/Population
439 Division; 2017.
- 440 16. **World Population Prospects: The 2017 Revision, Online Demographic Profiles**
441 [<https://population.un.org/wpp/Graphs/DemographicProfiles/>]
- 442 17. UNFPA, HAI: **Ageing in the Twenty-First Century: A celebration and a challenge.**
443 United Nations Population Fund (UNFPA), New York, and HelpAge International,
444 London; 2012.
- 445 18. **World Population Prospects: The 2010 Revision**
446 [<http://esa.un.org/unpd/wpp/unpp/p2k0data.asp>]
- 447 19. PRB: **2017 World Population Data Sheet** In. Washington, D.C: Population Reference
448 Bureau 2017.
- 449 20. UN: **Population Ageing and Development: Department of Economic and Social**
450 **Affairs, Population Division, United Nations.** 2009.
- 451 21. **AIDS by Numbers**
- 452 22. Linley L, Prejean J, An Q, Chen M, Hall HI: **Racial/ethnic disparities in HIV diagnoses**
453 **among persons aged 50 years and older in 37 US States, 2005-2008.** *American journal*
454 *of public health* 2012, **102**(8):1527-1534.
- 455 23. Chamie G, Clark TD, Kabami J, Kadede K, Ssemmondo E, Steinfeld R, Lavoy G,
456 Kwarisiima D, Sang N, Jain V *et al*: **A hybrid mobile approach for population-wide**
457 **HIV testing in rural east Africa: an observational study.** *The lancet HIV* 2016,
458 **3**(3):e111-e119.

- 459 24. **90–90–90 - An ambitious treatment target to help end the AIDS epidemic**
460 [<https://www.unaids.org/en/resources/documents/2017/90-90-90>]
- 461 25. **90-90-90: An ambitious treatment target to help end the AIDS epidemic**
462 [http://www.unaids.org/sites/default/files/media_asset/90-90-90_en.pdf]
- 463 26. Nabukenya AM, Matovu JKB: **Correlates of HIV status awareness among older**
464 **adults in Uganda: results from a nationally representative survey.** *BMC Public*
465 *Health* 2018, **18**(1):1128.
- 466 27. Hontelez JAC, de Vlas SJ, Baltussen R, Newell M-L, Bakker R, Tanser F, Lurie M,
467 Bärnighausen T: **The impact of antiretroviral treatment on the age composition of**
468 **the HIV epidemic in sub-Saharan Africa.** *AIDS (London, England)* 2012, **26**(0 1).
- 469 28. Mahy M, Autenrieth CS, Stanecki K, Wynd S: **Increasing trends in HIV prevalence**
470 **among people aged 50 years and older: evidence from estimates and survey data.**
471 *AIDS (London, England)* 2014, **28**(4):S453.
- 472 29. Justice A, Falutz J: **Ageing and HIV: an evolving understanding.** *Current opinion in*
473 *HIV and AIDS* 2014, **9**(4):291-293.
- 474 30. Brooks JT, Buchacz K, Gebo KA, Mermin J: **HIV infection and older Americans: the**
475 **public health perspective.** *American journal of public health* 2012, **102**(8):1516-1526.
- 476 31. Freeman E, Anglewicz P: **HIV prevalence and sexual behaviour at older ages in rural**
477 **Malawi.** *International journal of STD & AIDS* 2012, **23**(7):490-496.
- 478 32. Kharsany ABM, Karim QA: **HIV Infection and AIDS in Sub-Saharan Africa:**
479 **Current Status, Challenges and Opportunities.** *Open AIDS J* 2016, **10**:34-48.
- 480 33. PHIA: **Population-based HIV Impact Assessment (PHIA) survey.** In. New York,
481 USA: ICAP at Columbia University 2018.
- 482 34. Shisana O, Rehle T, Simbayi LC, Zuma K, Jooste S, Zungu N, Labadarios D, Onoya D:
483 **South African national HIV prevalence, incidence and behaviour survey, 2012.**
484 2014.
- 485 35. Schatz E, Knight L: **"I was referred from the other side": Gender and HIV testing**
486 **among older South Africans living with HIV.** *PloS one* 2018, **13**(4):e0196158-
487 e0196158.
- 488 36. Negin J, Gregson S, Eaton JW, Schur N, Takaruza A, Mason P, Nyamukapa C: **Rising**
489 **Levels of HIV Infection in Older Adults in Eastern Zimbabwe.** *PLoS One* 2016,
490 **11**(11):e0162967.
- 491 37. Mohlabane N, Tutshana B, Peltzer K, Mwisongo A: **Barriers and facilitators associated**
492 **with HIV testing uptake in South African health facilities offering HIV Counselling**
493 **and Testing.** *Health SA Gesondheid* 2016, **21**(1):86-95.
- 494 38. Negin J, Cumming RG: **HIV infection in older adults in sub-Saharan Africa:**
495 **extrapolating prevalence from existing data.** *Bulletin of the World Health*
496 *Organization* 2010, **88**:847-853.
- 497 39. Freeman E, Anglewicz P: **HIV prevalence and sexual behaviour at older ages in rural**
498 **Malawi.** *International journal of STD & AIDS* 2012, **23**(7):490-496.
- 499 40. Ford CL, Godette DC, Mulatu MS, Gaines TL: **Recent HIV Testing Prevalence,**
500 **Determinants, and Disparities Among U.S. Older Adult Respondents to the**
501 **Behavioral Risk Factor Surveillance System.** *Sexually transmitted diseases* 2015,
502 **42**(8):405-410.
- 503 41. Aday LA, Andersen RM: **A Framework for the Study of Access to Medical Care.**
504 *Health Services Research* 1974:208-220.

- 505 42. Andersen RM: **Revisiting the Behavioral Model and Access to Medical care: Does it**
506 **matter?** *Journal of Health and Social Behaviour* 1995, **36** (March):1-10.
- 507 43. Andersen RM: **National Health Surveys and the Behavioral Model of Health Services**
508 **Use.** *Med Care* 2008, **46**:647-653.
- 509 44. Negin J, Nemsler B, Cumming R, Lelera E, Amor YB, Pronyk P: **HIV attitudes,**
510 **awareness and testing among older adults in Africa.** *AIDS and Behavior* 2012,
511 **16**(1):63-68.
- 512 45. Akers A, Bernstein L, Henderson S, Doyle J, Corbie-Smith G: **Factors associated with**
513 **lack of interest in HIV testing in older at-risk women.** *Journal of Women's Health*
514 2007, **16**(6):842-858.
- 515 46. Oraka E, Mason S, Xia M: **Too old to test? Prevalence and correlates of HIV testing**
516 **among sexually active older adults.** *Journal of gerontological social work* 2018,
517 **61**(4):460-470.
- 518 47. Gwadz M, Leonard NR, Honig S, Freeman R, Kutnick A, Ritchie AS: **Doing battle with**
519 **"the monster:" how high-risk heterosexuals experience and successfully manage**
520 **HIV stigma as a barrier to HIV testing.** *International journal for equity in health* 2018,
521 **17**(1):46-46.
- 522 48. Madise NJ, Ziraba AK, Inungu J, Khamadi SA, Ezech A, Zulu EM, Kebaso J, Okoth V,
523 Mwau M: **Are slum dwellers at heightened risk of HIV infection than other urban**
524 **residents? Evidence from population-based HIV prevalence surveys in Kenya.**
525 *Health & place* 2012, **18**(5):1144-1152.
- 526 49. Gage AJ, Ali D: **Factors associated with self-reported HIV testing among men in**
527 **Uganda.** *AIDS Care* 2005, **17**(2):153-165.
- 528 50. UBOS, ICF: **Uganda Demographic and Health Survey 2016: Key Indicators Report.**
529 In. Kampala, Uganda and Rockville, Maryland, USA; 2017.
- 530 51. Scholten F, Mugisha J, Seeley J, Kinyanda E, Nakubukwa S, Kowal P, Naidoo N,
531 Boerma T, Chatterji S, Grosskurth H: **Health and functional status among older people**
532 **with HIV/AIDS in Uganda.** *BMC Public Health* 2011, **11**:886.
- 533 52. UBOS: **Uganda Population and Housing Census 2014.** In. Kampala, Uganda: Uganda
534 Bureau of Statistics; 2016.
- 535 53. **Masaka district** [<http://www.lcmt.org/uganda/masaka>]
536 54. **Hoima district** [<http://www.lcmt.org/uganda/masaka>]
- 537 55. Anderson TW, Kish L: **Survey Sampling.** In.; 1966.
- 538 56. UBOS, ICF International: **Uganda Demographic Health Survey 2016.** In. Kampala,
539 Uganda Uganda Bureau of Statistics (UBOS); 2017.
- 540 57. SurveyCto: **SurveyCTO: How it works.** In.; 2018.
- 541 58. Baeten JM, Donnell D, Kapiga SH, Ronald A, John-Stewart G, Inambao M, Manongi R,
542 Vwalika B, Celum C: **Male circumcision and risk of male-to-female HIV-1**
543 **transmission: a multinational prospective study in African HIV-1 serodiscordant**
544 **couples.** *AIDS (London, England)* 2010, **24**(5):737.
- 545 59. Gray RH, Kigozi G, Serwadda D, Makumbi F, Watya S, Nalugoda F, Kiwanuka N,
546 Moulton LH, Chaudhary MA, Chen MZ: **Male circumcision for HIV prevention in**
547 **men in Rakai, Uganda: a randomised trial.** *The Lancet* 2007, **369**(9562):657-666.
- 548 60. Kironde B, Wamala R, Kwagala B: **Determinants of male circumcision for HIV/AIDS**
549 **prevention in East Central Uganda.** *African journal of reproductive health* 2016,
550 **20**(1):80-87.

551 61. Negin J, Cumming RG: **HIV infection in older adults in sub-Saharan Africa:**
552 **extrapolating prevalence from existing data.** *Bulletin of the World Health*
553 *Organization* 2010, **88**(11):847-853.

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579 **Tables**

580 **Table 1 Descriptive characteristics of older persons in rural Uganda**

Variables	Number (n)	Percent (%)
-----------	------------	-------------

Age group		
50-59	312	48.3
60-69	176	27.2
70+	158	24.5
Sex of the respondents		
Female	334	51.5
Male	315	48.5
Education level		
None	111	17.1
Primary	378	58.2
Secondary or higher	160	24.7
Worked in the last 12 months		
No	308	47.5
Yes	341	52.5
Religion		
Catholic	279	43.0
Anglican	208	32.0
Muslim	99	15.3
Others	63	9.7
Currently in union		
No	262	40.4
Yes	387	59.6
Number of wives or husbands		
One	304	46.8
Two plus	83	12.8
Not in union	262	40.4
Correct knowledge about 4-5 HIV transmission statements		
No	230	35.4
Yes	419	64.6
Stigma on at least 4-8 statements		
No	436	67.2
Yes	213	32.8
Total	649	100.0

581

582

583

584

585

586

587

588

589

590 **Table 1 continued**

Variables	Number (n)	Percent (%)
Need to test for HIV once a year even when you know you are HIV negative		
No	35	5.4

Yes	614	94.6
Do older men who are 50 years older need to be circumcised to prevent HIV in		
No	202	31.1
Yes	447	68.9
Had sex in the last 12 months		
No	314	49.5
Yes	320	50.5
Number of lifetime sexual partners		
One	231	72.4
Two or more	88	27.6
Ever given or received money or gifts for sex		
No	490	75.5
Yes	159	24.5
Gave or received money or gifts for sex in the last 12 months		
No	574	88.4
Yes	75	11.6
Drinks alcohol		
No	426	65.6
Yes	223	34.4
Uses tobacco or drugs		
No	564	86.9
Yes	85	13.1
Self-reported STI in the last 12 months		
No	459	70.7
Yes	190	29.3
Male circumcised		
No	199	30.7
Yes	116	17.9
No, female	334	51.5
Ever tested for HIV and received results		
No	119	18.3
Yes	530	81.7
Tested for HIV and received results in the last 12 months		
No	307	47.3
Yes	342	52.7
Total	649	100

591

592

593

594 **Table 2 Association between recent HIV testing and background factors among older**595 **persons in Uganda**

**Tested for HIV and received results
in the last 12 months**

Variables	Yes (%)	Total (N)	P-value
Age group			<0.001
50-59	63.5	312	
60-69	50.6	176	
70+	34.2	158	
Sex of the respondents			0.21
Female	50.3	334	
Male	55.2	315	
Education level			0.10
None	44.1	111	
Primary	55.6	378	
Secondary or higher	51.9	160	
Work in the last 12 months			<0.001
No	45.8	308	
Yes	58.9	341	
Religion			0.09
Catholic	49.8	279	
Anglican	50	208	
Muslim	62.6	99	
Others	58.7	63	
Currently in union			<0.01
No	45.8	262	
Yes	57.4	387	
Number of wives or husbands			<0.01
One	54.9	304	
Two plus	66.3	83	
Not in union	45.8	262	
Correct Knowledge about 4-5 HIV transmission modes			0.01
No	46.1	230	
Yes	56.3	419	
Stigma on at least 4-8 statements			0.71
No	53.2	436	
Yes	51.6	213	
Test for HIV once a year even HIV negative			0.01
No	31.4	35	
Yes	53.9	614	
Total	52.7	100	

596

597

598 **Table 2 Continued**

Variables	Tested for HIV and received results in the last 12 months	
	Yes (%)	Total

Older men need to circumcise to prevent HIV			0.01
No	44.6	202	
Yes	56.4	447	
Had sex in the last 12 months			<0.001
No	38.9	314	
Yes	66.9	320	
Number of lifetime sexual partners			0.39
One	65.4	231	
Two or more	70.5	88	
Ever given or received money or gifts for sex			0.06
No	50.6	490	
Yes	59.1	159	
Gave or received money or gifts for sex in the last 12 months			0.01
No	50.9	574	
Yes	66.7	75	
Drinks alcohol			0.94
No	52.6	426	
Yes	52.9	223	
Uses tobacco or drugs			0.33
No	52	564	
Yes	57.6	85	
Self-reported STI in the last 12 months			<0.01
No	48.8	459	
Yes	62.1	190	
Male circumcised			<0.01
No	48.2	199	
Yes	67.2	116	
No, female	50.3	334	
Total	52.7	100	

599

600

601

602

603

604 **Table 3 Logistic regression of recent HIV testing against background factors, HIV knowledge and stigma and behavioural**
 605 **factors**
 606

Variables	Model 1 Adjusted Odds Ratios (aOR)	95% Confidence Interval	Model 2 Odds Ratios (aOR)	95% Confidence Interval	Model 3 Odds Ratios (aOR)	95% Confidence Interval
Age group						
50-59 (rc)	1				1.00	
60-69	0.61**	0.41-0.88	0.62*	0.43-0.92	0.78	0.52-1.18
70+	0.32***	0.22-0.49	0.36***	0.24-0.56	0.49**	0.31-0.79
Sex (rc = female)	1.08	0.75-1.55	1.11	0.77-1.59	0.79	0.47-1.13
Worked in the last 12 months (rc=No)	1.38*	1.00-1.93	1.38	1.01-2.10	1.34	0.96-1.91
Currently in union (rc = not in union)	1.46*	1.0102.09	1.46*	0.93-1.98	0.81	0.51-1.28
Correct knowledge on 4-5 HIV prevention modes			1.25	0.87-1.78	1.10	0.75-1.59
Agreement with on 4-8 stigma statements			1.03	0.72-1.47	1.01	0.69-1.45
Older men need to circumcise to prevent HIV (rc = No)			1.36	0.95-1.94	1.19	0.81-1.75
Self-reported sexually transmitted infections in the last 12 months (rc = No)					1.59*	1.09-2.30
Male circumcised						
No (rc)					1.00	
Yes					1.71*	1.00-2.93
Not Applicable to females					1.00	
Sexual activity in the last 12 months (rc= No)					2.89***	1.83-4.57
Transactional sex in the last 12 months (rc = No)					1.05	0.60-1.84
<i>Observations (N)</i>	646		646		631	

*aOR (Adjusted Odds Ratios); rc: reference category; 95% confidence intervals in second column: * p < 0.05, ** p < 0.01, *** p < 0.001;*

Model 1, controlling for socio-demographics, Model 2, controlling for HIV knowledge and stigma & Model 3, controlling for behavioural factors

607