

DISENTANGLING THE RISK OF MOTHERHOOD: AN EXPLORATORY STUDY OF MATERNAL MORTALITY IN INDIA

Background

The increased availability of household survey data and precise estimation techniques have enhanced our capabilities to understand maternal mortality. At present there are two estimation techniques which are commonly used to estimate maternal mortality through demographic and health surveys (DHS) in developing countries are sisterhood method (Graham et al., 1989) which is an indirect technique and second one is a direct variant of sisterhood method termed as direct sisterhood method (Rutenburg et al. 1991). Through these, we can have only a single index of maternal mortality. Apart from other studies which provide only a single estimate of maternal mortality, this study tried to estimate the mortality risk during and outside pregnancy period since, pregnancy-related risks can be better understood by studying the relationship between the mortality of pregnant women relative to the mortality of non-pregnant women (Stecklov, 1995). Moreover, general studies on maternal mortality in India gives overall estimates of maternal mortality ratio at the regional level and for different socio-economic subgroups but there are few studies done which made a comparison between the increased risk due to pregnancy for different socio-economic and demographic characteristics of Indian women. Since WHO states that 99% of all maternal deaths occur in developing countries and India still a significant contributor in the share, these maternal deaths are entirely preventable or avoidable, and literature which gives the precise estimates of avoidable mortality are scarcely available.

The methods which are currently are in use to estimate maternal mortality from different data sources worldwide are *Civil Registration (CR) and Vital Statistics Data, Census and population or household surveys, Sisterhood Methods, Reproductive Age Mortality Studies (RAMOS)*. Since there are different issues with the assumptions and estimations of these methods, this study uses a technique given by Stecklov in 1995 through which we can calculate separate mortality rates for women during and outside pregnancy period. By detaching pregnancy-related from non-pregnancy related mortality, will not only provide a measure of maternal mortality that is more coherent with the WHO definition of maternal mortality, but may also lead to new insights concerning the types of risks being measured (Stecklov, 1995). This study uses calendar data from the fourth round of the National Family Health Survey (2015-16), which is the Indian version of DHS to accomplish its objectives.

Research Questions

This study is based on two research questions which are as follows

1. What is the age pattern of mortality among women in reproductive ages during pregnancy period compared to non-pregnancy period?
2. Does this age pattern of mortality vary for different background characteristics?

Objective

To estimate mortality rates, and summary risk measures for India by different subpopulation during pregnancy and non-pregnancy period.

Data and Methods

The present study used secondary data from the fourth round of the National Family Health Survey (NFHS), which is the Indian version of the Demographic Health Survey. It is a large-scale, multi-round survey, which is conducted in a representative sample of households throughout India under the aegis of the Ministry of Health and Family Welfare, India. Fourth in the series, NFHS4, with some new additions was undertaken

in 2015-16. Unlike its previous three rounds, the fourth round National Family Health Survey encompasses and provides the estimates in the field of population, health, and nutrition to all the 29 States, 7 Union Territories, and 640 districts. With the prime objective of providing much-needed data in the area of health and family welfare and newly arisen affair in the related field. A total of 628,892 households (almost seven times as large as the household sample size in the first two rounds of NFHS) were covered, and 699686 ever-married women in the age group 15-49) were interviewed. The description of different methodology which have been used for our analysis is discussed below.


The methodology used to estimate Pregnancy period mortality rate (PPMR), Non-pregnancy period mortality rates (NPPMR) and Summary Measures

The numerator for PPMR and NPPMR

In the fourth round of National Family Health Survey number of women (who was a usual resident of the household) died during *childbirth or within two months after the end of pregnancy or childbirth* or outside this period is enumerated and here referred as ***death during pregnancy and non-pregnancy period*** respectively. Death during pregnancy and the non-pregnancy period are used as the numerator for the estimation of pregnancy and non-pregnancy period mortality rates (***PPMR***) & (***NPPMR***) for total females and females from selected background characteristics. Enumerated deaths in pregnancy and non-pregnancy period outside the age range of (15-49) are not considered for the calculation of rates. ***Deaths due to accidents, violence, poisoning, homicide or suicide are not included in the numerator.***

Denominator for PPMR and NPPMR

Denominator for the mortality rates which are person-years lived in pregnancy and non-pregnancy period' for the living and dead women in the age range (15-49). For the living women, 'person-years' lived in pregnancy and non-pregnancy period' is extracted through the calendar data which gives the month-wise history of women's pregnancy, birth and contraceptive usage up to 80 months. It provides the total number of months lived by women in the reference period of the study or total exposure. In the calendar of individual woman episodes of pregnancy, births and particular contraceptive usage which is basically months lived in pregnancy, birth months and the total number of months a particular contraceptive respectively is given. Months lived in pregnancy, month in which woman gave birth and the month in which pregnancy is terminated is recorded as "P," "B" and "T" respectively. The total number of "P", "B" and "T" is termed as months lived in pregnancy period or pregnancy period exposure. Non-pregnancy period for women is calculated by subtracting pregnancy period exposure from total exposure. Finally, all three exposure months is converted into years. On the other hand, no information is given about pregnancy period exposure in the reference period for the women who died during pregnancy period. So the pregnancy period exposure is simulated by assuming that

Pregnancy period exposure of dead women is similar as living women (e.g., pregnancy period exposure of a dead woman from the rural region is similar as pregnancy period exposure of living women from the rural region).  (Assumption 1)

The method to estimate PPMR, NPPMR and their summary measures is given by (Stecklov 1995), and here it is used with some modifications

The method to estimate PPMR, NPPMR and their summary measures is given by (Stecklov 1995), and here it is used with some modifications

$$PPMR = \frac{\text{Number of deaths in pregnancy period in the age group } (x, x + n)}{\text{Person years lived in pregnancy in the age group } (x, x + n)}$$

$$NPPMR = \frac{\text{Number of deaths in non – pregnancy period } (x, x + n)}{\text{Person years lived outside pregnancy in the age group } (x, x + n)}$$

Here non-pregnancy related mortality during the pregnancy period is assumed to equal non-pregnancy-related mortality during the non-pregnancy period. \longrightarrow (Assumption 2)

The assumption that the baseline (non-pregnancy-related) risks are constant regardless of whether a woman is in the pregnancy period appears to be a useful starting point.

Summary risk measures namely pregnancy to non-pregnancy mortality ratio, pregnancy-related mortality rate which gives attributable risk which can be attributed to pregnancy-related conditions, and pregnancy-related pregnancy period mortality which gives attributable risk% attributable to pregnancy-related conditions for an age interval (x, x+n) have been calculated with the help above two rates as follows,

$$\text{Pregnancy to non pregnancy mortality ratio} = \frac{PPMR}{NPPMR}$$

$$\text{Pregnancy related mortality rate} = (PPMR - NPPMR)$$

$$\text{Pregnancy related pregnancy period mortality}(\%) = \frac{(PPMR - NPPMR)}{PPMR}$$

Preliminary Findings-

In the age-group 30-49 mortality level during pregnancy period is approximately three times higher than the mortality levels in the non-pregnancy period. This age group also faces the highest proportionate risk from pregnancy-related mortality, with more than 75% mortality being pregnancy-related. These measures are lowest in the 15-19 age group. For all age groups combined, the data suggest that the mortality levels during pregnancy period are about two times higher than non-pregnancy pregnancy period, pregnancy to non-pregnancy mortality rate is 1.14 and 49% mortality in reproductive ages being pregnancy-related.

Age group	Pregnancy Period				Non-Pregnancy period				Summary risk measures		
	(A)	(B)	(C) ^a		(D)	(E)	(F) ^b		(G) ^c	(H) ^d	(I) ^e
	Deaths	Exposure	Mortality Rate	95% Confidence Limits	Deaths	Exposure	Mortality Rate	95% Confidence Limits	Pregnancy to non-pregnancy mortality ratio	Pregnancy-related mortality rate	Pregnancy-related pregnancy period mortality (%)
15-19	30	26,819	1.12	(0.72 , 1.52)	403	571,806	0.70	(0.64 , 0.77)	1.6	0.4	37.0
20-24	161	83,004	1.94	(1.64 , 2.24)	397	502,150	0.79	(0.71 , 0.87)	2.4	1.2	59.2
25-29	120	52,491	2.29	(1.88 , 2.70)	373	468,480	0.80	(0.72 , 0.88)	2.8	1.4	65.2
30-49	136	29,840	4.56	(3.79 , 5.32)	2263	1,355,191	1.67	(1.60 , 1.74)	2.7	2.9	63.4
Total	447	192,154	2.33	(2.11 , 2.54)	3436	2,897,627	1.19	(1.15 , 1.23)	1.9	1.1	49.0

^aA/B*1,000. ^bD/E*1,000. ^cC/F. ^dC-F. ^eH/C*100

The sensitivity of the Estimated Pregnancy and Non-Pregnancy Mortality Rates

The results of the sensitivity analysis of the mortality rates for total females to assumption 1 discuss is presented in figure 3.1 and 3.2. Three separate measures of pregnancy mortality are shown in figure 3.1. The total height of each bar is showing the estimate of maternal mortality which can be similar to the estimates that we can have from applying direct sisterhood method. The method used here suggests that the total height of each bar which is pregnancy-period mortality rate can be divided into two parts. The first part is baseline mortality which is the bottom part of each bar and the second part is pregnancy-related mortality which upper part of each bar. In figure 3.1, the mortality rates that are shown are estimated with taking an exposure of dead women in the denominator which is extracted with the help of assumption 1. According to assumption 1, the pregnancy exposure of dead women is same as living women. It means that the proportion of pregnancy exposure out of total exposure for the living women in the reference period is the same for the women who died in the reference period. However, in figure 3.2 denominator of the mortality rates shown do not contain the exposure of dead women. After comparative assessment of both figures, it is evident that the age pattern of mortality rates is similar and there is no such significant change occurred in the rates if the exposure of dead women is not considered. It seems that

since the portion of exposure of dead women in the denominator is too smaller than living women, so its effect is getting diluted. In conclusion, we can say that assumption 1 will not hamper our process of estimating mortality rates and their summary measures, and we can rely on estimated mortality rates.




Figure 3.1: Pregnancy and Non-pregnancy related mortality rates with exposure of dead women in the denominator, by age-group, India, 2011-16




Figure 3.2: Pregnancy and Non-pregnancy related mortality rates without exposure of dead women in the denominator, by age-group, India, 2011-16

References

- Radkar, A., & Parasuraman, S. (2007). Maternal deaths in India: an exploration. *Economic and Political Weekly*, 3259-3263.
- Alkema, L., Chou, D., Hogan, D., Zhang, S., Moller, A. B., Gemmill, A., ... & Say, L. (2016). Global, regional, and national levels and trends in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *The Lancet*, 387(10017), 462-474.
- RGI (various years) Special bulletin on maternal mortality in India. New Delhi: *Office of the Registrar General & Census Commissioner, India, New Delhi*
- Government of India (2005) National Rural Health Mission (2005–2012): Mission Document. *Ministry of Health and Family Welfare, Government of India, New Delhi, p. 28*
- Srinivasan, K., Chander, S. & Arokiasamy, P. (2007) Reviewing reproductive and child health programmes in India. *Economic & Political Weekly* 42(27), 2931–2939.
- Jain, A. K. (2010). Janani Suraksha Yojana and the maternal mortality ratio. *Economic and Political Weekly*, 45(11), 15-16.
- Tanner, J. C., AM, A. R., Candland, T., Galdo, V., Manang, F., Trichler, R., & Yamagata, R. (2014). Delivering the Millennium Development Goals to reduce maternal and child mortality: a systematic review of impact evaluation evidence.
- Jeffrey, P and R Jeffrey (2010) Only when the boat has started sinking: A maternal death in rural north India. *Social Science and Medicine*, 71: 1711–18